



INTERNATIONAL CENTRE
OF EXCELLENCE FOR
EDUCATION IN
MATHEMATICS

The Improving Mathematics Education in Schools (TIMES) Project

INTRODUCTION TO COORDINATE GEOMETRY

A guide for teachers - Years 9–10

NUMBER AND ALGEBRA ■
Module 29

June 2011

YEARS
9
10

Introduction to Coordinate Geometry

(Number and Algebra : Module 29)

For teachers of Primary and Secondary Mathematics

510

Cover design, Layout design and Typesetting by Claire Ho

The Improving Mathematics Education in Schools (TIMES) Project 2009-2011 was funded by the Australian Government Department of Education, Employment and Workplace Relations.

The views expressed here are those of the author and do not necessarily represent the views of the Australian Government Department of Education, Employment and Workplace Relations.

© The University of Melbourne on behalf of the International Centre of Excellence for Education in Mathematics (ICE-EM), the education division of the Australian Mathematical Sciences Institute (AMSI), 2010 (except where otherwise indicated). This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. 2011.

<http://creativecommons.org/licenses/by-nc-nd/3.0/>



The Improving Mathematics Education in Schools (TIMES) Project

INTRODUCTION TO COORDINATE GEOMETRY

A guide for teachers - Years 9–10

NUMBER AND ALGEBRA •
Module 29

June 2011

Peter Brown
Michael Evans
David Hunt
Janine McIntosh
Bill Pender
Jacqui Ramagge

YEARS
9
10

INTRODUCTION TO COORDINATE GEOMETRY

ASSUMED KNOWLEDGE

- Fluency with the arithmetic of the rational numbers
- Knowledge of ratios
- Congruent and similar triangles
- Basic algebraic notation
- Fluency with algebraic expressions and equations
- Basic plotting points in the Cartesian plane including plotting points from a table of values.

MOTIVATION

Coordinate geometry is one of the most important and exciting ideas of mathematics. In particular it is central to the mathematics students meet at school. It provides a connection between algebra and geometry through graphs of lines and curves. This enables geometric problems to be solved algebraically and provides geometric insights into algebra.

The invention of calculus was an extremely important development in mathematics that enabled mathematicians and physicists to model the real world in ways that was previously impossible. It brought together nearly all of algebra and geometry using the coordinate plane. The invention of calculus depended on the development of coordinate geometry.


CONTENT

It is expected that students have met plotting points on the plane and have plotted points from tables of values of both linear and non linear functions.

The number plane (Cartesian plane) is divided into four quadrants by two perpendicular axes called the **x-axis** (horizontal line) and the **y-axis** (vertical line). These axes intersect at a point called the **origin**. The position of any point in the plane can be represented by an **ordered pair** of numbers (x, y) . These ordered pairs are called the coordinates of the point.

The point with coordinates $(4, 2)$ has been plotted on the Cartesian plane shown. The coordinates of the origin are $(0, 0)$.

Once the coordinates of two points are known the distance between the two points and midpoint of the interval joining the points can be found.



DISTANCE BETWEEN TWO POINTS

Distances are always positive, or zero if the points coincide. The distance from A to B is the same as the distance from B to A . We first find the distance between two points that are either vertically or horizontally aligned.

EXAMPLE

Find the distance between the following pairs of points.


a $A(1, 2)$ and $B(4, 2)$

b $A(1, -2)$ and $B(1, 3)$

SOLUTION


a The distance $AB = 4 - 1 = 3$

Note: The distance AB is obtained from the difference of the x -coordinates of the two points.




b The distance $AB = 3 - (-2) = 5$

Note: The distance AB is obtained from the difference of the y -coordinates of the two points.



The example above considered the special cases when the line interval AB is either horizontal or vertical. Pythagoras' theorem is used to calculate the distance between two points when the line interval between them is neither vertical nor horizontal.

The distance between the points $A(1, 2)$ and $B(4, 6)$ is calculated below.



$$AC = 4 - 1 = 3 \text{ and } BC = 6 - 2 = 4.$$


By Pythagoras' theorem,

$$AB^2 = 3^2 + 4^2 = 25$$

And so $AB = 5$

The general case

We can obtain a formula for the length of any interval. Suppose that $P(x_1, y_1)$ and $Q(x_2, y_2)$ are two points.



Form the right-angled triangle PQX , where X is the point (x_2, y_1) ,

$$PX = x_2 - x_1 \text{ or } x_1 - x_2 \quad \text{and} \quad QX = y_2 - y_1 \text{ or } y_1 - y_2$$

depending on the positions of P and Q .

By Pythagoras' theorem:

$$\begin{aligned} PQ^2 &= PX^2 + QX^2 \\ &= (x_2 - x_1)^2 + (y_2 - y_1)^2 \end{aligned}$$

Therefore $PQ = QP = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Note that $(x_2 - x_1)^2$ is the same as $(x_1 - x_2)^2$ and therefore it doesn't matter whether we go from P to Q or from Q to P – the result is the same.

EXAMPLE

Find the distance between the points $A(-4, -3)$ and $B(5, 7)$.

SOLUTION

In this case, $x_1 = -4$, $x_2 = 5$, $y_1 = -3$ and $y_2 = 7$.

$$\begin{aligned} AB^2 &= (x_2 - x_1)^2 + (y_2 - y_1)^2 \\ &= (5 - (-4))^2 + (7 - (-3))^2 \\ &= 9^2 + 10^2 \\ &= 181 \end{aligned}$$

Thus, $AB = \sqrt{181}$

Note that we could have chosen $x_1 = 5$, $x_2 = -4$, $y_1 = 7$ and $y_2 = -3$ and still obtained the same result. As long as (x_1, y_1) refers to one point and (x_2, y_2) the other point, it does not matter which one is which.

EXERCISE 1

Show that the distance between the points $A(a, b)$ and $B(c, d)$ is the same as the distance between

- the points $P(a, d)$ and $Q(c, b)$
- the points $U(b, a)$ and $V(d, c)$

Illustrate both of these.

EXERCISE 2

The distance between the points $(1, a)$ and $(4, 8)$ is 5. Find the possible values of a and use a diagram to illustrate.

THE MIDPOINT OF AN INTERVAL

The coordinates of the midpoint of a line interval can be found using averages as we will see.

We first deal with the situation where the points are horizontally or vertically aligned.

EXAMPLE

Find the coordinates of the midpoint of the line interval AB , given:


a $A(1, 2)$ and $B(7, 2)$

b $A(1, -2)$ and $B(1, 3)$

SOLUTION


- a AB is a horizontal line interval, the midpoint is at $(4, 2)$, since 4 is halfway between 1 and 7.

Note: 4 is the average of 1 and 7, that is, $4 = \frac{1+7}{2}$.




- b The midpoint of AB has coordinates $(1, \frac{1}{2})$.

Note that $\frac{1}{2}$ is the average of 3 and -2 .



When the interval is not parallel to one of the axes we take the average of the x-coordinate and the y-coordinate. This is proved below.



Let M be the midpoint of the line AB . Triangles AMS and MBT are congruent triangles (AAS), and so $AS = MT$ and $MS = BT$.

Hence the x-coordinate of M is the average of 1 and 5.

$$x = \frac{5+1}{2} = 3$$


The y coordinate of M is the average of 2 and 8.

$$y = \frac{2+8}{2} = 5$$

Thus the coordinates of the midpoint M are $(3, 5)$.

The general case

We can find a formula for the midpoint of any interval. Suppose that $P(x_1, y_1)$ and $Q(x_2, y_2)$ are two points and let $M(x, y)$ be the midpoint.



Triangles PMS and MQT are congruent triangles (AAS), and so $PS = MT$ and $MS = QT$.

Hence the x -coordinate of M is the average of x_1 and x_2 , and y -coordinate of M is the average of y_1 and y_2 . Therefore

$$x = \frac{x_1 + x_2}{2} \quad \text{and} \quad y = \frac{y_1 + y_2}{2}$$

Midpoint of an interval

The midpoint of an interval with endpoints $P(x_1, y_1)$ and $Q(x_2, y_2)$ is $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$.

Take the average of the x -coordinates and the average of the y -coordinates.

EXAMPLE

Find the coordinates of the midpoint of the line interval joining the points $(6, 8)$ and $(-3, 2)$.

SOLUTION

The midpoint has coordinates, $\left(\frac{6 + (-3)}{2}, \frac{8 + 2}{2}\right) = \left(\frac{3}{2}, 5\right)$

EXAMPLE

If $C(3, 6)$ is the midpoint of line interval AB and A has coordinates $(-1, 1)$, find the coordinates of B .

SOLUTION

Let the coordinates of B be (x_1, y_1) .

$$\frac{x_1 + (-1)}{2} = 3 \quad \text{and} \quad \frac{y_1 + 1}{2} = 6$$

$$x_1 - 1 = 6 \quad y_1 + 1 = 12$$

so $x_1 = 7$ so $y_1 = 11$.

Thus B has coordinates $(7, 11)$.

EXERCISE 3


A square has vertices $O(0, 0)$, $A(a, 0)$, $B(a, a)$ and $C(0, a)$.

- Find the midpoint of the diagonals OB and CA .
- Find the length of a diagonal of the square and the radius of the circle in which $OABC$ is inscribed.
- Find the equation of the circle inscribing the square.

THE GRADIENT OF A LINE

Gradient of an interval

The gradient is a measure of the steepness of line. There are several ways to measure steepness. In coordinate geometry the standard way to define the gradient of an interval AB is $\frac{\text{rise}}{\text{run}}$ where **rise** is the change in the y -values as you move from A to B and **run** is the change in the x -values as you move from A to B . We will usually use the pronumeral m for gradient.



Given the points $A(2, 1)$ and $B(5, 6)$:

$$\text{gradient of interval } AB = \frac{\text{rise}}{\text{run}} = \frac{6-1}{5-2} = \frac{5}{3}$$


Notice that as you move from A to B along the interval the y -value increases as the x -value increases. The gradient is **positive**.

Given the points $A(2, 7)$ and $B(6, 1)$

$$\text{gradient of interval } AB = \frac{\text{rise}}{\text{run}} = \frac{1-7}{6-2} = \frac{-6}{4} = -\frac{3}{2}$$

or

$$\text{gradient of interval } BA = \frac{\text{rise}}{\text{run}} = \frac{7-1}{2-6} = -\frac{3}{2}$$




Notice that in this case as we move from A to B the y value decreases as the x value increases. The gradient is **negative**. Similarly the gradient of $BA = -\frac{3}{2}$ which is the same as the gradient of AB .

In general:


gradient of line interval $AB = \frac{\text{rise}}{\text{run}}$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$



Note that since $\frac{y_2 - y_1}{x_2 - x_1} = \frac{y_1 - y_2}{x_1 - x_2}$ it does not matter which point we take as the first and which point we take as the second.

If the interval is vertical, the run is zero and the gradient of the interval is not defined. This is shown by interval AB .




Gradient of PQ is zero
Gradient of AB is not defined

If the interval is horizontal, the rise is zero as shown by interval PQ . The gradient of the interval is zero.

Gradient of a line

The gradient of a line is defined to be the gradient of any interval within the line.

This definition depends on the fact that two intervals on a line have the same gradient.



Suppose AB and PQ are two intervals on the same straight line. Draw right-angled triangles ABX and PQY with sides AX and PY parallel to the x -axis and sides BX and QY parallel to the y -axis.

Triangle ABX is similar to triangle PQY since the corresponding angles are equal.

Therefore:

$$\frac{QY}{PY} = \frac{BX}{AX}$$


That is, the intervals have the same gradient.

EXAMPLE

A line passes through the points (1, 2) and (5, 10). Find its gradient.

SOLUTION

$$\begin{aligned} \text{gradient} &= \frac{y_2 - y_1}{x_2 - x_1} \\ &= \frac{10 - 2}{5 - 1} \\ &= 2 \end{aligned}$$




EXAMPLE

A line passes through the point (5, 7) and has gradient $\frac{2}{3}$. Find the x-coordinate of a point on the line when $y = 13$.

SOLUTION

Gradient of the line = $\frac{6}{x-5}$. Thus,

$$\begin{aligned} \frac{6}{x-5} &= \frac{2}{3} \\ 18 &= 2(x-5) \\ 9 &= x-5 \\ x &= 14 \end{aligned}$$




EXERCISE 4

Find the gradient of the line passing through (a, b) and (0, c)

Intercepts

The **x-intercept** of a line is the point at which it crosses the x-axis.

The **y-intercept** of a line is the point at which it crosses the y-axis.



In the diagram to the left the y-intercept is at A and the x-intercept at B.


The second diagram shows a line parallel to the x -axis and it has a y -intercept at C .

The third diagram shows a line parallel to the y -axis and it has an x -intercept at D .

EQUATION OF A STRAIGHT LINE

When we plot points which satisfy the equation $y = 2x + 1$ we find that they lie in a straight line.

Can we find the equation of the line given suitable geometric information about the line? The following shows that this can be done given the gradient of the line and the y -intercept.



THE LINE $y = 3x + 2$


Consider the line with gradient 3 and y -intercept 2. This passes through the point $A(0, 2)$. Let $B(x, y)$ be any point on this line.

$$\begin{aligned} \text{Gradient of interval } AB &= \frac{\text{rise}}{\text{run}} \\ &= \frac{y-2}{x-0} \\ &= \frac{y-2}{x} \end{aligned}$$

The gradient of the line is 3.

So,
$$\frac{y-2}{x} = 3$$

Rearranging
$$\begin{aligned} y - 2 &= 3x \\ y &= 3x + 2 \end{aligned}$$



So the coordinates of $B(x, y)$ satisfies $y = 3x + 2$. This is called the equation of the line.

Conversely suppose that $B(x, y)$ satisfies the equation $y = 3x + 2$, then $\frac{y-2}{x-0} = 3$ and it passes the point $(0, 2)$ so the point lies on the line with gradient 3 and y -intercept 2.


We summarise this by saying that the equation of the line is $y = 3x + 2$.

THE EQUATION $y = mx + c$

Consider the line with gradient m and y -intercept c . If passes through the point $A(0, c)$. Let $B(x, y)$ be any point on this line.

$$\begin{aligned} \text{Gradient of interval } AB &= \frac{y-c}{x-0} \\ &= \frac{y-c}{x} \end{aligned}$$

We know the gradient of the line is m .




$$\begin{aligned}\text{Therefore } \frac{y-c}{x} &= m \\ y-c &= mx \\ y &= mx + c\end{aligned}$$

That is, the line in the cartesian plane with gradient m and y -intercept c has equation $y = mx + c$. Conversely, the points whose coordinates satisfy the equation $y = mx + c$ always lie on the line with gradient m and y -intercept c .

Vertical and horizontal lines

Vertical lines

In a vertical line all points have the same x -coordinate, but the y -coordinate can take any value. The equation of the vertical line through the point $(6, 0)$ is $x = 6$. The x -axis intercept is 6. All the points on this line have x -coordinate 6.




In general, the equation of the vertical line through $P(a, b)$ is $x = a$.

Because this line does not have a gradient it cannot be written in the form

$$y = mx + b.$$

Horizontal lines

A horizontal line has gradient 0. In a horizontal line all points have the same y -coordinate, but the x -coordinate can take any value. The equation of the horizontal line through the point $(0, 5)$ is $y = 5$. The equation of the horizontal line through the point $(9, 5)$ is $y = 5$.



In general, the equation of the horizontal line through $P(a, b)$ is $y = b$.

EXAMPLE

Write down the gradient and y -intercept of the line with equation $y = 3x - 4$.

SOLUTION

The gradient of the line is 3 and the y -intercept is -4 .

Sometimes an equation needs to be rearranged before the gradient and y -intercept can be determined. Consider the following example.

EXAMPLE

Rewrite the equation $2x + 3y = 6$ in the form $y = mx + c$ and hence find the value of the gradient and y -intercept.

SOLUTION

$$2x + 3y = 6$$

so $3y = 6 - 2x$

that is, $y = 2 - \frac{2x}{3}$

Thus $y = -\frac{2x}{3} + 2$

The gradient of the line is $-\frac{2}{3}$ and the y -intercept is 2.


Equation of a line given its gradient and a point on the line

We want to find the equation of the line with gradient m and which passes through the point $P(x_1, y_1)$.

Let $P(x, y)$ be any point with $x \neq x_1$ on the line passing through the point $A(x_1, y_1)$ and let m be the gradient of this line.

Using gradient, $m = \frac{y - y_1}{x - x_1}$

and $y - y_1 = m(x - x_1)$



This is the equation of the straight line with gradient m passing through the point $A(x_1, y_1)$.

EXAMPLE

Find the equation of the line that passes through the point $(-2, 3)$ with gradient -4 .

SOLUTION

The equation for this line is:

$$y - y_1 = m(x - x_1)$$

$$y - 3 = -4(x - (-2))$$

that is,
$$\begin{aligned} y - 3 &= -4x - 8 \\ y &= -4x - 5 \end{aligned}$$

Note that it is usual to give the answer in the form $y = mx + c$

Equation of a straight line given two points

Given two points $A(x_1, y_1)$ and $B(x_2, y_2)$ the equation of the line passing through the two points can be found.

The gradient m of the line passing through $A(x_1, y_1)$ and $B(x_2, y_2) = \frac{y_2 - y_1}{x_2 - x_1}$, $x_2 \neq x_1$

Substituting into $y - y_1 = m(x - x_1)$ gives

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1), \quad x_2 \neq x_1$$

GRAPHING STRAIGHT LINES

Using the equation to sketch the line

If you are given an equation of a straight line and asked to draw its graph all you need to do is find two points whose coordinates satisfy the equation and plot the points. There are two commonly used methods to find two points.

Using the y-intercept and one other point

Using the y-intercept and a second point the equation can be found .

EXAMPLE

Draw the graph of $y = 2x + 3$.


SOLUTION

The y-intercept is 3 and the gradient 2.

Substitute $x = 1$, so $y = 5$ giving the point $(1, 5)$ lies on the line.

Plot the two points and draw the line through them.

This method does not work if the line is parallel to the y-axis.



EXAMPLE

The gradient of a line is -6 and the y -intercept is 2 . Find the equation of the line and sketch it.

SOLUTION


Using the $y = mx + c$ form for the equation of a straight line.

The equation of the line is $y = -6x + 2$.

The point $(0, 2)$ lies on the line.

$$\begin{aligned} \text{Substitute } x = 1 \text{ in } y &= -6x + 2 \\ &= -6 \times 1 + 2 \\ &= -4 \end{aligned}$$

The point $(1, -4)$ lies on the line. The graph is shown.

**x-intercept y-intercept method**

In this method both intercepts are found.

- The x -intercept is found by substituting $y = 0$ and
- The y -intercept is found by substituting $x = 0$.

This method does not work if the line is parallel to an axis or passes through the origin.

EXAMPLE


Using the x -intercept y -intercept method sketch the graph of:

a $y = 3x - 4$

b $2x + 3y + 6 = 0$


SOLUTION

a When $x = 0$, $y = -4$
 When $y = 0$, $3x - 4 = 0$
 $3x = 4$
 $x = \frac{4}{3}$



b When $x = 0$, $3y + 6 = 0$
 $3y = -6$
 $y = -2$

When $y = 0$, $2x + 6 = 0$
 $2x = -6$
 $x = -3$



The general form for the equation of a straight line

The equations $y = 2x - 3$, $x = 6$ and $2x - 3y = 6$ can be written as $-2x + y + 3$, $x - 6 = 0$ and $2x - 3y - 6 = 0$ respectively.

The **general** form for the equation of a line is $ax + by + c = 0$ where a , b and c are constants and $a \neq 0$ or $b \neq 0$. The equation of every line can be put in general form. The general form is not unique. The equation $x + 2y + 1 = 0$ is the same straight line as $2x + 4y + 2 = 0$.

EXERCISE 5

An equilateral triangle ABC has coordinates $O(0, 0)$, $B(a, 0)$ and $C(c, d)$.


- Find c and d in terms of a by using the fact that $OB = BC = CO$.
- Find the equation of the lines OB , BC and CO .

PARALLEL AND PERPENDICULAR LINES

Parallel lines

If two lines l_1 and l_2 are parallel then corresponding angles are equal.

Conversely, if corresponding angles are equal then the lines are parallel.




Theorem

Two lines are **parallel** if they have the same gradient and conversely, two lines with the same gradient are parallel.

Proof

In the diagram, two lines are drawn and the right-angled triangles PQX and ABY are added with $QX = BY$.



If the lines are parallel then $\angle PQX = \angle ABY$ (corresponding angles).

The two triangles are congruent by the AAS test.

Therefore $PX = PY$ and $\frac{PX}{QX} = \frac{AY}{BY}$.

That is, the gradients are equal.

Conversely. If the gradients are equal $\frac{PX}{QX} = \frac{AY}{BY}$.

Now $QX = BY$ and therefore $PX = AY$.

Hence the triangles QPX and ABY are congruent by the SAS test.

Hence the corresponding angles PQX and ABY are equal and the lines are parallel.

EXAMPLE

Show that the line passing through the points $A(6, 4)$ and $B(7, 11)$ is parallel to the line passing through $P(0, 0)$ and $Q(2, 14)$.

SOLUTION

$$\text{Gradient of } AB = \frac{11-4}{7-6} = 7$$

$$\text{Gradient of } PQ = \frac{14-0}{2-0} = 7$$

The two lines have the same gradient and so are parallel.

EXAMPLE

Find the equation of the line that is parallel to the line $y = -2x + 6$ and passing through the point $A(1, 10)$.

SOLUTION

The gradient of the line $y = -2x + 6$ is -2 .

Therefore the line through the point $A(1, 10)$ parallel to $y = -2x + 6$ has equation:

$$y - y_1 = m(x - x_1)$$

$$y - 10 = -2(x - 1)$$

$$y = -2x + 12$$


Perpendicular lines

When we draw $y = 3x$ and draw a line perpendicular to it passing through the origin then it is clear that $y = -ax$ where a is a small positive number. We will show that the equation is $y = -\frac{1}{3}x$.

We are now going to show the surprising result that 'if two lines are **perpendicular** then the product of their gradients is -1 ' (or if one is vertical and the other horizontal). The converse is also true. That is, 'If the product of the gradients of two lines is -1 then they are perpendicular'.

We first consider the case when both lines pass through the origin.

Draw two lines passing through the origin with one of the lines having positive gradient and the other negative gradient.




Form right-angled triangles OPQ and OAB with $OQ = OB$.

$$\text{Gradient of the line } OA = \frac{AB}{OB}$$

$$\text{Gradient of the line through } OP = -\frac{OQ}{PQ}$$

$$\begin{aligned} \text{The product} &= -\frac{OQ}{PQ} \times \frac{AB}{OB} \\ &= -\frac{OQ}{PQ} \times \frac{AB}{OQ} \quad (\text{since } OQ = OB) \\ &= -\frac{AB}{PQ} \end{aligned}$$

If the lines are perpendicular, $\angle POQ = \angle AOB$.



Therefore $\triangle OPQ \equiv \triangle OAB$ (AAS),

so $PQ = AB$ and the product, $-\frac{AB}{PQ}$, of the gradients is -1 .

Conversely If the product is -1 , then $AB = PQ$,

so $\triangle OAB \equiv \triangle OPQ$ (SAS).

Therefore $\angle POQ = \angle AOB$ and so $\angle AOP = 90^\circ$.

We have now proved the result for lines through the origin.

If we are given two lines anywhere in the plane, we can draw lines through the origin parallel to the given two lines. The gradient of each new line is the same as the gradient of the corresponding original line. So the result holds for lines that do not necessarily pass through the origin.

EXAMPLE

Show that the line through the points $A(6, 0)$ and $B(0, 12)$ is perpendicular to the line through $P(8, 10)$ and $Q(4, 8)$.

SOLUTION

$$\begin{aligned} \text{Gradient of } AB &= \frac{12-0}{0-6} \\ &= -2 \end{aligned}$$

$$\begin{aligned} \text{Gradient of } PQ &= \frac{10-8}{8-4} \\ &= \frac{2}{4} \\ &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} (\text{Gradient of } AB) \times (\text{gradient of } PQ) &= -2 \times \frac{1}{2} \\ &= -1 \end{aligned}$$

Hence the lines are perpendicular.

Parallel and perpendicular lines

- If two non-vertical lines are **parallel** then they have the same gradient.
Conversely if two non-vertical lines have the same gradient then they are parallel.
- If two non-vertical lines are **perpendicular** then the product of their gradients is -1 . Conversely if the product of the gradients of two lines is -1 then they are perpendicular.

EXAMPLE

Find the equation of the line which passes through the point (1, 3) and is perpendicular to the line whose equation is $y = 2x + 1$.

SOLUTION

Gradient of the line $y = 2x + 1$ is 2.

Gradient of a line perpendicular to this line is $-\frac{1}{2}$.

The required equation is $y - 3 = -\frac{1}{2}(x - 1)$

$$2(y - 3) = -(x - 1)$$

$$2y + x = 7$$

Thus the equation of the required line is $2y + x = 7$.

PROOFS WITH COORDINATE GEOMETRY

Coordinate geometry can be used to prove results in Euclidean Geometry. An important aspect of doing this is placing objects on the Cartesian plane in a way that minimises calculations.

EXAMPLE

Prove that the midpoints of a parallelogram bisect each other using coordinate geometry.

SOLUTION

Let the coordinates of the vertices be $O(0, 0)$, $A(a, 0)$, $B(a + c, d)$ and $C(c, d)$.

There is no loss in generality in placing the vertices of the parallelogram on the Cartesian plane in this way.

The midpoint M of $OB = \left(\frac{a+c}{2}, \frac{d}{2}\right)$

The midpoint N of $AC = \left(\frac{a+c}{2}, \frac{d}{2}\right)$

$M = N$ and so the midpoints coincided which means that the diagonals bisect each other.

EXAMPLE

Prove that the diagonals of a rhombus bisect each other at right angles using coordinate geometry.

SOLUTION

Let the coordinates of the vertices be $O(0, 0)$, $A(a, 0)$, $B(a + c, d)$ and $C(c, d)$.

Because it is a rhombus all the sides are of equal length.

$$OA = AB = BC = CO$$

$$\text{Gradient of } OB = \frac{d}{a+c} \text{ and,}$$

$$\text{Gradient of } AC = \frac{d}{c-a}$$

$$\text{The product of the gradients of the diagonals} = \frac{d}{a+c} \times \frac{d}{c-a} = \frac{d^2}{c^2-a^2}.$$

$$OA^2 = a^2 \text{ and by Pythagoras' theorem, } AB^2 = c^2 + d^2$$

$$\text{so } a^2 = c^2 + d^2.$$

$$\text{Hence } d^2 = -(c^2 - a^2)$$

Thus the product of the gradients of the diagonals = -1 .

EXERCISE 6

In any triangle ABC prove that $AB^2 + AC^2 = 2(AD^2 + DC^2)$

Where D is the midpoint of BC .

EXERCISE 7

Prove that set of points equidistant from two given points is a straight line.

EXERCISE 8

Prove that the lines joining the midpoints of opposite sides of a quadrilateral and the lines joining the midpoints of its diagonals meet in a point and bisect each other.

LINKS FORWARD

Coordinate geometry leads into many other topics in school mathematics. The techniques of coordinate geometry are used in calculus, functions, statistics and many other important areas.

HISTORY

There were three facets of the development of coordinate geometry.

- The invention of a system of coordinates
- The recognition of the correspondence between geometry and algebra
- The graphic representation of relations and functions

The Greek mathematician Menaechmus (380–320 BC) proved theorems by using a method that was very close to using coordinates and it has sometimes been maintained that he had introduced coordinate geometry.



Apollonius of Perga (262–190 BC) dealt with problems in a manner that may be called an coordinate geometry of one dimension; with the question of finding points on a line that were in a ratio to the others. The results and ideas of the ancient Greeks provided a motivation for the development of coordinate geometry.

Coordinate geometry has traditionally been attributed to René Descartes (1599–1650) and Pierre de Fermat (1601–1665) who independently provided the beginning of the subject as we know it today.

ANSWERS TO EXERCISES


EXERCISE 1

$$AB^2 = PQ^2 = UV^2 = (d - b)^2 + (c - a)^2$$



EXERCISE 2

$$a = 4 \text{ or } a = 12$$



EXERCISE 3

a $\left(\frac{a}{2}, \frac{a}{2}\right)$

b diagonal = $\sqrt{2}a$, radius = $\frac{\sqrt{2}a}{2}$

c $\left(x - \frac{a}{2}\right)^2 + \left(y - \frac{a}{2}\right)^2 = \frac{a^2}{2}$

EXERCISE 4

$$\frac{b-c}{a}$$

EXERCISE 5

$$\mathbf{a} \quad c = \frac{a}{2}, d = \frac{\sqrt{3}a}{2} \quad \mathbf{b} \quad OA: y = 0, OC: y = \frac{d}{c}, OB: y = \frac{d}{c-a}(x-a)$$

EXERCISE 6

Place the triangle so D is at the origin. Then let the coordinates of B and C be $(-a, 0)$ and $(a, 0)$ respectively. Let the coordinates of A be (d, c) .

$$AB^2 = c^2 + (d+a)^2 \text{ and } AC^2 = c^2 + (d-a)^2$$

$$\text{So } AB^2 + AC^2 = 2c^2 + 2d^2 + 2a^2$$

$$AD^2 = c^2 + d^2 \text{ and } DC^2 = a^2$$

$$\text{Hence } AB^2 + AC^2 = 2(AD^2 + DC^2)$$

EXERCISE 7

Let $P(x, y)$ be a point equidistant from $A(a, b)$ and $C(c, d)$

$$PA = PC$$

$$2(c-a)x + 2(d-b)y = d^2 + c^2 - a^2 - b^2$$

EXERCISE 8

Let the coordinates of the vertices be $O(0, 0)$, $A(a, c)$, $B(m, n)$ and $D(b, 0)$

Show that the midpoint of all the required line segments has coordinates

$$\left(\frac{1}{4}(m+b+a), \frac{1}{4}(n+c)\right)$$



INTERNATIONAL CENTRE
OF EXCELLENCE FOR
EDUCATION IN
MATHEMATICS

The aim of the International Centre of Excellence for Education in Mathematics (ICE-EM) is to strengthen education in the mathematical sciences at all levels—from school to advanced research and contemporary applications in industry and commerce.

ICE-EM is the education division of the Australian Mathematical Sciences Institute, a consortium of 27 university mathematics departments, CSIRO Mathematical and Information Sciences, the Australian Bureau of Statistics, the Australian Mathematical Society and the Australian Mathematics Trust.



AUSTRALIAN MATHEMATICS TRUST



The ICE-EM modules are part of *The Improving Mathematics Education in Schools (TIMES) Project*.

The modules are organised under the strand titles of the Australian Curriculum:

- Number and Algebra
- Measurement and Geometry
- Statistics and Probability

The modules are written for teachers. Each module contains a discussion of a component of the mathematics curriculum up to the end of Year 10.

www.amsi.org.au